

Static and Dynamical properties of environment-induced topological phase transitions

Fabrizio Pavan¹, G. De Filippis^{1*}

1. University of Naples "Federico II", Dept. of Physics "E. Pancini", via Cinthia, 21, 80126 Napoli, Italy

* Contributing authors within the NQSTI partnership; others are listed in the reference [4]

Abstract: We study environmental effects on the Su–Schrieffer–Heeger model with local baths, showing that suitable couplings can induce topological transitions that are detectable using both static and dynamical observables.

We study the nonperturbative effects of environmental coupling on the Su-Schrieffer-Heeger (SSH) chain, where local harmonic oscillator baths interact with either intracell or intercell transfer integrals. Contrary to common assumptions, this type of coupling, when properly engineered, can induce transitions to topological phases. To move beyond the weak coupling regime described by Lindbladian formalism, we use a world-line quantum Monte Carlo (QMC) technique [1] to map the full phase diagram of the model (see Fig. 1 (a) and (b)). In this framework, we identify the probability distribution of polarization as a key marker of topological phase transitions under open boundary conditions. Notably, this distribution exhibits a bimodal behaviour in the topological phase, both with and without environmental interactions, highlighting the emergence of edge states (see Fig. 1 (c)). Additionally, we provide a qualitative and analytical description using cluster perturbation theory (CPT) [3], which is well-suited for lattice models with dominant local interactions under periodic boundary conditions. This formalism enables accurate calculations of dressed Green's functions, allowing us to determine spectral functions and the topological invariant [2]. The discontinuous behaviour of the topological invariant reveals the critical coupling strength at which the topological phase transition occurs, in agreement with Monte Carlo results. This critical point is further marked by the closing of the energy gap between the two bands, observed as the merging of the two main peaks in the spectral functions. Moreover, we demonstrate that environmental interactions introduce non-Hermitian effects in the fermionic subsystem [4], providing an effective non-Hermitian Hamiltonian description. Finally, to demonstrate that this effect can be experimentally probed through the Mean Chiral Displacement (MCD) [5], we simulate the interacting single-electron dynamics using the Tensor Network formalism [6].




Fig. 1 . Phase diagram, in the case of intra[inter]cell hopping coupled with local baths in (a) [(b)], for quantum (red diamonds) and classical bosons (grey points) through MC technique. Green line denotes the critical values of the couplings obtained through CPT for quantum baths. (c) Polarization probability distribution in the case of intercell hopping coupled with local baths.

References:

- [1] H. G. Evertz, The loop algorithm, *Adv. Phys.* 52, 1 (2003).
- [2] Z. Wang and S.-C. Zhang, Strongly correlated topological superconductors and topological phase transitions via Green's function, *Phys. Rev. B* 86, 165116 (2012).
- [3] D. Sénéchal, D. Perez, and M. Pioro-Ladrière, Spectral weight of the Hubbard model through cluster perturbation theory, *Phys. Rev. Lett.* 84, 522 (2000).
- [4] F. Pavan, A. de Candia, G. Di Bello, V. Cataudella, N. Nagaosa, C. A. Perroni, and G. De Filippis, Witnessing environment induced topological phase transitions via quantum Monte Carlo and cluster perturbation theory studies, *Phys. Rev. B* 109, 115147 (2024).
- [5] F. Cardano et al., Detection of Zak phases and topological invariants in a chiral quantum walk of twisted photons, *Nature Communications* 8, 15516 (2017)
- [6] F. Pavan, G. Di Bello, G. De Filippis, C. A. Perroni, Single particle dynamical signature of topology induced by single mode cavities in Su-Schrieffer-Heeger chain, arXiv:2512.09520.

